

ABSTRACTING LYMPHOMA CANCER INCIDENCE AND TREATMENT DATA
February 8, 2007

Quiz 1

1. Which of the following cluster of differentiation is typical of B-cells?
 - a. CD 4
 - b. CD 8
 - c. CD20
 - d. None of the above
2. A bone marrow biopsy performed 1/1/2007 is positive for B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma (B-CLL/SLL). An excisional biopsy of a cervical lymph node performed one week later confirms this diagnosis. What is the ICD-O 3 histology code?
 - a. 8140/3 Adenocarcinoma
 - b. 9670/3 Small lymphocytic lymphoma
 - c. 9823/3 Chronic lymphocytic leukemia
 - d. 8000/3 Malignancy NOS
3. A patient presents with lymphoma involving supraclavicular (C77.0) and pretracheal (C77.0) lymph nodes. What is the ICD-O 3 topography code?
 - a. C77.9
 - b. C77.8
 - c. C80.9
 - d. C77.0
4. Which of the following are types of cutaneous T-cell lymphoma? (You may pick more than one)
 - a. Bowens Disease
 - b. Mycosis Fungoides
 - c. Sezary Syndrome
 - d. Sarcomatoid Carcinoma
5. A patient with a history of Waldenstroms macroglobulinemia (9761/3) now presents with follicular lymphoma, grade 2 (9691/3).

According to the SEER "Single versus Subsequent Primaries of Lymphatic and Hematopoietic Diseases" table this is one primary.

- a. True
- b. False

Quiz 2

1. Which of the following is an extralymphatic site?
 - a. Lymph Nodes
 - b. Stomach
 - c. Spleen
 - d. Thymus

2. A patient presents with lymphoma. Staging work-up indicates bilateral supraclavicular lymph node involvement. No further disease identified. What stage is this?
 - a. I
 - b. II
 - c. III
 - d. IV

3. A patient is found to have lymphoma of the right hilar lymph nodes with direct extension to the right lung. No further disease is identified. What Stage is this?
 - a. Stage I
 - b. Stage Ie
 - c. Stage Iie
 - d. Stage IV

4. A single lesion in which of the following sites would indicate stage IV lymphoma?
 - a. Axillary lymph nodes
 - b. Stomach
 - c. Thymus
 - d. Liver

5. A patient was presents with lymphoma. Staging work-up shows nodal involvement above and below the diaphragm with involvement of the spleen, what is the extension code for Collaborative Stage?
 - a. 12
 - b. 30
 - c. 31
 - d. 32

Quiz 3

1. Patient with widespread abdominal and inguinal lymphadenopathy has two inguinal lymph nodes excised. They are found to be positive for lymphoma. How do we code the excision of these lymph nodes?
 - a. 02 Diagnostic Staging Procedure
 - b. 15 Surgery of Primary Site
 - c. 25 Surgery of Primary Site
 - d. 99 Surgery of Primary Site
2. A patient with a lymphoma originating in the stomach presents for a total gastrectomy and lymph node dissection. 3 of 7 pyloric lymph nodes were positive for lymphoma. What is Scope of Regional Lymph Node Surgery? (see slides for codes)
 - a. 0
 - b. 3
 - c. 4
 - d. 5
3. External beam radiation applied to the mantle field is targeting lymph nodes:
 - a. Above the diaphragm
 - b. In the abdomen
 - c. In the inguinal area
 - d. In the brain
4. _____ is a drug that works by binding to a specific protein (the CD20 antigen) on the surface of B-cells. From there, it works with the body's own immune system to attack and kill the marked B-cells.
 - a. Adriamycin
 - b. Vincristine
 - c. Rituximab (Rituxan)
 - d. Etoposide
5. A lymphoma patient presents for a peripheral blood stem cell harvest and high dose chemotherapy. The stem cells are harvested 1/1/07. Chemotherapy begins 1/15/07 and the stem cell cells are reintroduced one 2/1/07. If this was the only treatment given, what would be the Date Systemic Therapy Started?
 - a. 1/1/07
 - b. 1/15/07
 - c. 2/1/07
 - d. 99/99/07